



Large Artery Stiffening and Remodeling Are Independently Associated With All-Cause Mortality and Cardiovascular Events in Chronic Kidney Disease

Submitted by Emmanuel Lemoine on Thu, 03/26/2015 - 14:24

Titre	Large Artery Stiffening and Remodeling Are Independently Associated With All-Cause Mortality and Cardiovascular Events in Chronic Kidney Disease
Type de publication	Article de revue
Auteur	Karras, Alexandre [1], Haymann, Jean-Philippe [2], Bozec, Erwan [3], Metzger, Marie [4], Jacquot, Christian [5], Maruani, Gerard [6], Houillier, Pascal [7], Froissart, Marc [8], Stengel, Bénédicte [9], Guardiola, Philippe [10], Laurent, Stéphane [11], Boutouyrie, Pierre [12], Briet, Marie [13]
Editeur	American Heart Association
Type	Article scientifique dans une revue à comité de lecture
Année	2012
Langue	Anglais
Date	2012/01/12
Numéro	6
Pagination	1451 - 1457
Volume	60
Titre de la revue	Hypertension
ISSN	1524-4563
Mots-clés	aortic stiffness [14], carotid artery common [15], Mortality [16], Remodeling [17], Renal Insufficiency [18]

Résumé en anglais	<p>Chronic kidney disease, even at moderate stages, is characterized by a high incidence of cardiovascular events. Subclinical damage to large arteries, such as increased arterial stiffness and outward remodeling, is a classical hallmark of patients with chronic kidney disease. Whether large artery stiffness and remodeling influence the occurrence of cardiovascular events and the mortality of patients with chronic kidney disease (stages 2–5) is still debated. This prospective study included 439 patients with chronic kidney disease (mean age, 59.8±14.5 years) with a mean measured glomerular filtration rate of 37 mL/min per 1.73 m². Baseline aortic stiffness was estimated through carotid-femoral pulse wave velocity measurements; carotid stiffness, diameter, and intima-media thickness were measured with a high-resolution echotracking system. For the overall group of patients, the 5-year estimated survival and cumulative incidence of cardiovascular events were 87% and 16%, respectively. In regression analyses adjusted on classical cardiovascular and renal risk factors, aortic stiffness remained significantly associated with all-cause mortality (for 1 SD, Cox model-derived relative risk [95% CI], 1.48 [1.09–2.02]) and with fatal and nonfatal cardiovascular events (for 1 SD, Fine and Gray competing risks model-derived relative risk [95% CI], 1.35 [1.05–1.75]). Net reclassification improvement index was significant (29.0% [2.3–42.0%]). Carotid internal diameter was also independently associated with all-cause mortality. This study shows that increased aortic stiffness and carotid internal diameter are independent predictors of mortality in patients with stages 2 to 5 chronic kidney disease and that aortic stiffness improves the prediction of the risk.</p>
URL de la notice	http://okina.univ-angers.fr/publications/ua9188 [19]
DOI	10.1161/HYPERTENSIONAHA.112.197210 [20]
Lien vers le document	http://dx.doi.org/10.1161/HYPERTENSIONAHA.112.197210 [20]

Liens

- [1] [http://okina.univ-angers.fr/publications?f\[author\]=16415](http://okina.univ-angers.fr/publications?f[author]=16415)
- [2] [http://okina.univ-angers.fr/publications?f\[author\]=16416](http://okina.univ-angers.fr/publications?f[author]=16416)
- [3] [http://okina.univ-angers.fr/publications?f\[author\]=16417](http://okina.univ-angers.fr/publications?f[author]=16417)
- [4] [http://okina.univ-angers.fr/publications?f\[author\]=16418](http://okina.univ-angers.fr/publications?f[author]=16418)
- [5] [http://okina.univ-angers.fr/publications?f\[author\]=16419](http://okina.univ-angers.fr/publications?f[author]=16419)
- [6] [http://okina.univ-angers.fr/publications?f\[author\]=16420](http://okina.univ-angers.fr/publications?f[author]=16420)
- [7] [http://okina.univ-angers.fr/publications?f\[author\]=16421](http://okina.univ-angers.fr/publications?f[author]=16421)
- [8] [http://okina.univ-angers.fr/publications?f\[author\]=16422](http://okina.univ-angers.fr/publications?f[author]=16422)
- [9] [http://okina.univ-angers.fr/publications?f\[author\]=16423](http://okina.univ-angers.fr/publications?f[author]=16423)
- [10] <http://okina.univ-angers.fr/philippe.guardiola/publications>
- [11] [http://okina.univ-angers.fr/publications?f\[author\]=16424](http://okina.univ-angers.fr/publications?f[author]=16424)
- [12] [http://okina.univ-angers.fr/publications?f\[author\]=16425](http://okina.univ-angers.fr/publications?f[author]=16425)
- [13] [http://okina.univ-angers.fr/publications?f\[author\]=13896](http://okina.univ-angers.fr/publications?f[author]=13896)
- [14] [http://okina.univ-angers.fr/publications?f\[keyword\]=14855](http://okina.univ-angers.fr/publications?f[keyword]=14855)
- [15] [http://okina.univ-angers.fr/publications?f\[keyword\]=14856](http://okina.univ-angers.fr/publications?f[keyword]=14856)
- [16] [http://okina.univ-angers.fr/publications?f\[keyword\]=7726](http://okina.univ-angers.fr/publications?f[keyword]=7726)
- [17] [http://okina.univ-angers.fr/publications?f\[keyword\]=7143](http://okina.univ-angers.fr/publications?f[keyword]=7143)
- [18] [http://okina.univ-angers.fr/publications?f\[keyword\]=7678](http://okina.univ-angers.fr/publications?f[keyword]=7678)
- [19] <http://okina.univ-angers.fr/publications/ua9188>
- [20] <http://dx.doi.org/10.1161/HYPERTENSIONAHA.112.197210>